

Brain–behavior relationships in the experience and regulation of negative emotion in healthy children: Implications for risk for childhood depression

DAVID PAGLIACCIO, JOAN L. LUBY, KATHERINE R. LUKING, ANDREW C. BELDEN, AND DEANNA M. BARCH

Washington University in St. Louis

Abstract

Structural and functional alterations in a variety of brain regions have been associated with depression and risk for depression across the life span. A majority of these regions are associated with emotion reactivity and/or regulation. However, it is generally unclear what mechanistic role these alterations play in the etiology of depression. A first step toward understanding this is to characterize the relationships between variation in brain structure/function and individual differences in depression severity and related processes, particularly emotion regulation. To this end, the current study examines how brain structure and function predict concurrent and longitudinal measures of depression symptomatology and emotion regulation skills in psychiatrically healthy school-age children ($N = 60$). Specifically, we found that smaller hippocampus volumes and greater responses to sad faces in emotion reactivity regions predict increased depressive symptoms at the time of scan, whereas larger amygdala volumes, smaller insula volumes, and greater responses in emotion reactivity regions predict decreased emotion regulation skills. In addition, larger insula volumes predict improvements in emotion regulation skills even after accounting for emotion regulation at the time of scan. Understanding brain–behavior relationships in psychiatrically healthy samples, especially early in development, will help inform normative developmental trajectories and neural alterations in depression and other affective pathology.

Major depressive disorder (MDD) is a potentially devastating illness with a high lifetime prevalence, yet the available treatments are inadequate or ineffective for a large portion of those affected (Kessler, Berglund, & Demler, 2003). Furthermore, MDD has a high and rising prevalence during adolescence (Burke, Burke, Rae, & Regier, 1991). It is also evident in school-age children (Birmaher, Ryan, Williamson, & Brent, 1996; Kaufman, Martin, King, & Charney, 2001) and even in preschool-age children, where preschool-onset depression shows homotypic continuity in early childhood and strongly predicts DSM-V MDD in later childhood and early adolescence (Luby, Si, Belden, Tandon, & Spitznagel, 2009). These findings underscore the need for work identifying depression

risk factors early in development that can guide early identification and intervention, with aims to both treat early-onset depression and to ameliorate negative trajectories that may be set into place early in development.

Much foundational work has been done in adult populations to characterize core deficits associated with MDD and their neural underpinnings. This work is integral to identifying key systems disrupted in depression; however, to address goals of early intervention and prevention, research must be extended earlier in development in order to explore the relationship between individual differences in these core deficits and their putative neural mechanisms in healthy individuals. Here, we focus specifically on emotion processing, because adult and pediatric MDD have been related to psychological and behavioral differences in emotion processing as well as alterations in emotion processing neural circuitry.

We will investigate how structural and functional disruptions in several key regions of relevance to emotion reactivity and/or regulation predict individual differences in depressive symptomatology and emotion regulation skills in psychiatrically healthy children, with both cross-sectional and longitudinal analyses. Below, we briefly review work identifying two broad systems in the brain relevant to emotion reactivity and emotion regulation and examine evidence for their functional impairment in depression. We also review evidence for changes in the structure of several specific regions within these networks in depression, with a particular focus on the

This work was supported by the National Institute of Mental Health Grants MH64769 (to J.L.L.) and MH090786 (to J.L.L., D.M.B., and K.N.B.), a NARSAD award (to J.L.L.), and by Grant 1K01MH090515-01 from the National Institute of Mental Health (to A.C.B.). We thank all participants and their families who provided time and effort to make this study possible. The National Institute of Mental Health had no further role in the design and conduct of the study (collection, management, analysis, or interpretation of the data) or in the preparation, review, or approval of the manuscript. The first author had full access to all study data and takes responsibility for the integrity of the data and accuracy of the data analysis. The authors have no biomedical financial interests or potential conflicts of interest.

Address correspondence and reprint requests to: David Pagliaccio, Program in Neuroscience, Washington University in St. Louis, Campus Box 1125, One Brookings Drive, St. Louis, MO 63130; E-mail: david.pagliaccio@wustl.edu or david.pagliaccio@gmail.com.

insula, hippocampus, and amygdala, each of which is thought to play a key role in emotion reactivity and/or regulation.

Emotion reactivity and regulation networks

Task-based functional magnetic imaging (fMRI) studies in healthy adults suggest that emotion reactivity and regulation activate several neural systems with both common and distinct components (McRae et al., 2012; Ochsner, Bunge, Gross, & Gabrieli, 2002; Ochsner et al., 2004, 2009; Silvers et al., 2012). In particular, a “bottom-up” system, involved in the generation of emotion, has been shown to include the amygdala, striatum, hippocampus, and temporal regions that typically show increased reactivity to emotionally evocative stimuli (Ochsner & Feldman Barrett, 2001; Ochsner & Gross, 2005, 2007, 2008; Ochsner & Phelps, 2007). It is important that these regions show functional responses to emotionally evocative stimuli in children and adolescents as well, though the specific patterns of activity in response to stimulus valence may vary across age (e.g., Guyer et al., 2008; Monk et al., 2003; Pagliaccio et al., 2013; Thomas et al., 2001).

In contrast, studies of emotion regulation have consistently highlighted important roles for dorsal regions of the prefrontal cortex and the parietal cortex in a network of “top-down” regions that implement control processes for modulating and regulating activity in the limbic and subcortical systems (Buhle et al., 2013; Diekhof, Geier, Falkai, & Gruber, 2011; Ochsner, Bunge, Gross, & Gabrieli, 2002, 2004, 2009; Silvers et al., 2012). Consistent with these findings in adults, school-age children also show recruitment of top-down regions during reappraisal of negative emotional content (Belden, Luby, Pagliaccio, & Barch, 2014; McRae et al., 2012). It should also be noted that activity in top-down control regions during regulation may increase with age (presumably as regulation efficacy increases), while some research suggests that self-reported negative affect and amygdala reactivity may not show the same changes with age (McRae et al., 2012).

In concert with the above task-based fMRI studies, studies of resting-state functional connectivity in healthy adults have also supported this distinction between brain regions involved in the experience of emotion and those involved in the regulation of emotion. Briefly, resting-state functional connectivity represents correlations between spontaneous fluctuations in regional activity measured when a participant is at rest; this type of correlation may be interpreted as the “background pattern” of communication between brain regions reflecting a history of co-activation. Connectivity studies in adults have shown positive correlations between resting-state fluctuations in amygdala activity and fluctuations in activity in regions such as the hippocampus, striatum, and temporal lobe regions (Anticevic, Repovs, & Barch, 2010; Roy et al., 2009; Stein et al., 2007), that is, similar to the bottom-up network described above. In contrast, a host of frontal/parietal control regions show negative correlations

with fluctuations in amygdala activity (Anticevic et al., 2010; Roy et al., 2009; Stein et al., 2007), that is, similar to the top-down network. Both task and resting-state studies suggest that positive connectivity between the amygdala and bottom-up regions like the hippocampus remain relatively stable over development, while connectivity between the amygdala and top-down cortical regions strengthens over development (Gabard-Durnam et al., 2014; Vink, Derks, Hoogendam, Hillegers, & Kahn, 2014). Studies suggest that patients with depression may exhibit alterations in functional connectivity between the amygdala, cingulate, and insula (Connolly et al., 2013; Matthews, Strigo, & Simmons, 2008) and between prefrontal cortex regions and the cingulate and hippocampus (Greicius et al., 2007; Sheline, Price, Yan, & Mintun, 2010). In addition, childhood depression and maternal depression have been associated with reductions in connectivity between the amygdala and both these top-down and bottom-up regions (Luking et al., 2011), potentially relating to disruptions in both emotion processing and regulation abilities.

Amygdala

As noted above, there is a network of regions in the brain that respond to emotionally evocative stimuli. Among these regions, the amygdala has perhaps received the most attention. More specifically, functional activity of the amygdala in response to emotional stimuli is one of the most well-studied neural markers in affective psychopathology. The amygdala responds to highly emotionally salient stimuli of either positive or negative emotional valence (Davidson & Irwin, 1999). While the amygdala responds to both positive and negative emotional stimuli, it is generally found to show greater responses to emotional versus neutral stimuli (Fusar-Poli et al., 2009). Yet most studies to date have focused on the role of the amygdala in processing negative emotional stimuli, particularly threat- and sadness-related images. Along this line, it has been established that stronger amygdala responses to negative emotional faces are observed with increasing negative affect (Davidson & Irwin, 1999). Amygdala reactivity to emotional faces is increased in adolescents as compared to adults, which may relate to the increased vulnerability to affective disorders during adolescence (Guyer et al., 2008). Further suggestive of important developmental differences, there is evidence that differential activation to different face types may not be evident in school-age children, who show undifferentiated amygdala reactivity to a variety of emotional and neutral face types (Pagliaccio et al., 2013).

Nonetheless, depression has been related to amygdala hyperreactivity to negative emotional stimuli in adults (Groenewold, Opmeer, de Jonge, Aleman, & Costafreda, 2012; Hamilton et al., 2012; Stuhrmann, Suslow, & Dannlowski, 2011) and in children and adolescents (e.g., Barch, Gaffrey, Botteron, Belden, & Luby, 2012; Beesdo et al., 2009; Gaffrey et al., 2011; Pagliaccio et al., 2011; Yang et al., 2010). Specifically, preschool-onset depression severity has been shown

to correlate with increasing amygdala responses to sad versus neutral faces in school-age children (Barch et al., 2012; Gaffrey et al., 2011). Further, children/adolescents at risk for depression based on a parental history of MDD show elevated amygdala reactivity to fearful faces (Monk et al., 2008). Limbic hyperactivity may also be clinically relevant because it has been associated with treatment outcomes (Canli et al., 2005; Siegle, Carter, & Thase, 2006) and this hyperactivity may remit with successful antidepressant treatment (Fu et al., 2004).

Despite the focus that the amygdala has received in task-based fMRI studies, there have been relatively few investigations of amygdala volume in depressed and high-risk samples. One impediment is that the amygdala is a small structure with less distinct boundaries, making the measurement of volume more difficult to conduct reliably. Furthermore, studies that have investigated amygdala volumes have produced mixed results, with some reporting larger volumes (Frodl et al., 2002, 2004; Lange & Irle, 1999; Saleh et al., 2012; Van Eijndhoven et al., 2009; Vassilopoulou et al., 2012) and others reporting smaller amygdala volumes in depressed groups (Hastings, Parsey, Oquendo, Arango, & Mann, 2004; Sheline, Gado, & Price, 1998; Sheline, Sanghavi, Mintun, & Gado, 1999; Siegle, Konecky, Thase, & Carter, 2003). In pediatric populations, risk and protective factors have been related to amygdala volume, with some evidence that larger volumes are related to poorer outcomes. For example, early life stress (late adoption after institutional rearing) predicts larger amygdala volumes later in childhood, which in turn has been associated with anxiety and internalizing symptoms (Tottenham et al., 2010). Similarly, larger amygdala volumes have been related to greater fearfulness among girls, especially those with a family history of MDD (van der Plas, Boes, Wemmie, Tranel, & Nopoulos, 2010). Conversely, positive parenting during childhood predicted attenuated growth of amygdala volume into adolescence (Whittle et al., 2013).

Hippocampus

Although not often the focus of studies on emotional reactivity, the hippocampus has been shown to be active during the processing of facial emotion (Gur et al., 2002; Williams et al., 2001), where its response may correlate with increasing intensities of fearful face expression (Surguladze et al., 2003). Developmental differences in hippocampal reactivity are also evident, such that hippocampal responses to face stimuli strengthen across pubertal development (Moore et al., 2012). Positive functional connectivity between the hippocampus and amygdala may strengthen from adolescence to adulthood (Guyer et al., 2008). Structure-function correlations suggest that smaller hippocampal volumes are related to greater cortico-limbic responses to negative emotional faces in children (Suzuki et al., 2012). Examining depression, hippocampal responses to sad faces tend to be elevated among depressed patients; this hyperreactivity did not change with antidepressant treatment (Fu et al., 2004). In school-age

children, depression symptom severity at preschool age positively predicted hippocampal response to emotional faces (Barch et al., 2012). However, it is currently unclear whether alterations in hippocampal responses to emotion are a risk factor for depression, a result of depression, and/or indicative of a specific underlying mechanistic alteration in depression (e.g., increased reactivity to emotional content).

The hippocampus plays a critical role in the regulation of the stress response (Jacobson & Sapolsky, 1991), and there is consistent evidence that hippocampal volumes are altered by experiences of adversity and stress (e.g., Carballedo et al., 2012; Luby et al., 2013; Stein, Koverola, Hanna, Torchia, & McClarty, 1997; Woon & Hedges, 2008). Given the role of early life stress in the etiology of depression (Green et al., 2010; Kessler & Magee, 2009) and evidence for altered stress reactivity in depression (e.g., Burke, Davis, Otte, & Mohr, 2005; Heuser, Yassouridis, & Holsboer, 1994; Morris, Rao, & Garber, 2012), a number of studies have examined hippocampal volumes in depression. These studies have consistently shown reduced bilateral hippocampal volume in adult patients with depression (for meta-analyses, see Campbell & MacQueen, 2004; Videbech & Ravnkilde, 2004). This has been suggested as a potential risk factor for depression because volume reductions are observed in girls at risk for depression based on parental history (Amico et al., 2011; Chen, Hamilton, & Gotlib, 2010). One hypothesis is that early life stress may be a key causal factor in hippocampal volume reductions (e.g., Dannlowski et al., 2012). Alternatively, volume reductions may be a marker of deleterious effects of MDD, because the severity of volume reductions tends to correlate with the number of MDD episodes experienced (Videbech & Ravnkilde, 2004). In addition, smaller hippocampal volumes may predict poorer clinical outcomes with antidepressant treatment, while treatment may also contribute to increases in volume over time (Frodl et al., 2008).

Despite robust findings in adult depression, the literature on changes in hippocampal volume in childhood depression is mixed, with some reporting decreased volume (e.g., Caetano et al., 2007; MacMaster & Kusumakar, 2004; MacMaster et al., 2008; Rao et al., 2010) and others failing to detect an effect of childhood depression (e.g., Rosso et al., 2005), of exposure to maternal depression (Lupien et al., 2011), or of early life stress (Tottenham et al., 2010). This may be consistent with animal literature suggesting that negative factors, like early life stress, show a delayed effect on hippocampal volume that is not evident until later in development (Andersen & Teicher, 2004). However, other findings suggest that examining individual differences in risk factors for depression may reveal early hippocampal reductions; for example, poverty has been related to hippocampal reductions in children, as mediated by increased adverse parenting and life stress (Luby et al., 2013).

Insula

The insular cortex is a key region implicated in emotion processing. In particular, it has been suggested that the anterior

insula, in particular, may be crucial for the experience of emotion and interoceptive awareness (Gu, Hof, Friston, & Fan, 2013; Park & Tallon-Baudry, 2014). Further, the anterior insula may be a key waypoint for integrating top-down regulatory signals with bottom-up information about interoceptive experiences (Gu et al., 2013). Additional work suggests that the anterior insula may be involved in social emotions, such as guilt (Lamm & Singer, 2010). Recent work suggests that the anterior insula may be important for integrating information about the outcome of events in the world with internal goals and context factors to help support the generation of complex feeling states, which may include guilt (Koban & Pourtois, 2014). Consistent with this hypothesis, meta-analytic work shows robust activation of the insula in response to face stimuli displaying angry or disgusted expressions (Fusar-Poli et al., 2009). Insula activity has been consistently shown to be reduced in MDD patients in resting PET/SPECT studies and in response to negative stimuli in task-based fMRI studies (Fitzgerald, Laird, Maller, & Daskalakis, 2008). Further, in structural MRI studies, anterior insula gray matter volume has been shown to be reduced in people with current or remitted depression (Liu et al., 2014). Insula gray matter volume and density have also been shown to be reduced in first-episode MDD patients (Lai & Wu, 2014; Peng et al., 2011). Furthermore, volumetric changes in the anterior insula may predict treatment response in adults (Mayberg et al., 2000) and may serve as a biomarker for chronicity in early-onset MDD (Belden et al., 2014).

Summary and Goals for the Current Study

Given this literature identifying regions involved in emotion processing that show alterations in structure and function in depression and in high-risk states, we sought to explore relationships between these factors and depressive symptomology and emotion regulation skills in psychiatrically healthy school-age children. Specifically, we tested the hypotheses that depression-related alterations in both the structure and the function of brain regions involved in emotion reactivity and regulation would predict greater subclinical depressive symptomology and decreased ability to regulate sadness in 8- to 12-year-olds. Furthermore, we tested whether these neural factors would predict change in depressive symptomology and sadness regulation skills over approximately 18 months. Examining neural function, we took a broad network perspective and examined reactivity to sad faces in two networks of brain regions thought to be involved in emotion reactivity, including the amygdala, hippocampus, and additional medial temporal regions, or thought to be involved in emotion regulation, including the dorsal frontal and parietal regions (Luking et al., 2011). We hypothesized that greater responses to sad faces compared to neutral faces in the emotion reactivity-related network would predict greater depressive symptomology and poorer sadness regulation skills at baseline and would predict worsening of these outcomes over time. For the emotion regulation-related regions, we

hypothesized that reduced activity would predict greater depressive symptomology and poorer sadness regulation skills at baseline and would predict worsening of these outcomes over time. We also tested whether volume of the bilateral amygdala, hippocampus, and insula would predict these outcomes. For both the hippocampus and the insula, based on the literature linking depression and reduced volumes in adults, we hypothesized that smaller volumes would be associated with greater depressive symptomology and poorer sadness regulation skills at baseline and would predict worsening of these outcomes 18 months later. As noted above, the literature on amygdala volume in depression is mixed, but given findings that negative factors tend to predict larger volumes, we predicted that larger amygdala volumes would predict greater depressive symptomology and poorer sadness regulation skills at baseline and would predict worsening of these outcomes 18 months later.

Methods

Participants

The participants were a subsample of the children enrolled in the Preschool Depression Study (PDS), a prospective longitudinal investigation of preschoolers and their families conducted in the Early Emotional Development Program at the Washington University School of Medicine (WUSM). The current study reports on 60 children from the PDS (of the 305 total enrolled in the study) who did not have any prior history of a psychiatric disorder or any current disorders at the time of scan and who did not have a history of head trauma, neurological disease, or developmental delay. Forty-eight of these children had longitudinal data at an 18-month follow-up (see Table 1). Participants were between the ages of 8 and 12 ($M = 9.8$, $SD = 1.3$) at the time of the first scan and between the ages of 9 and 14 ($M = 11.2$, $SD = 1.2$) at the time of the follow-up. Parental written consent and child assent were obtained prior to participation and the institutional review board at WUSM approved all experimental procedures.

Recruitment for the PDS study occurred in two waves. The first wave recruited 3- to 6-year-old preschoolers from sites throughout the Saint Louis area for participation in a study examining the nosology of preschool depression. Recruitment was conducted through primary care practices and preschools/daycares that were accessible to the general community to ensure the socioeconomic and ethnic diversity of the sample. Recruitment sites were chosen at random using a geographically stratified method. The aim of this sampling technique was to recruit a large group of depressed preschoolers as well as smaller groups of disruptive and healthy preschoolers for comparison. To achieve this goal, caregivers completed a validated screening checklist, the Preschool Feelings Checklist (PFC; Luby, Heffelfinger, Koenig-McNaught, Brown, & Spitznagel, 2004). Previous studies indicated that a PFC score of 3 or higher has high sensitivity and specificity for diagnosing depression (Luby et al., 2004). In addition to

Table 1. Demographic and clinical characteristics of participants

	Baseline Scan (<i>N</i> = 60)			18-Month Follow-Up (<i>N</i> = 48)		
	Mean	<i>SD</i>	Range	Mean	<i>SD</i>	Range
Age (years)	9.78	1.25	8–12	11.2	1.22	9–14
Sex (% males)	55			56		
Ethnicity (% Caucasian)*	58			50		
Parental education	2.70	0.93	1–4	2.75	0.89	1–4
Parental income	2.92	1.15	1–4	3.08	1.03	1–4
CDI-C	40.53	4.82	35–57	39.81	7.4	34–83
CEMS sadness coping subscale	10.8	1.97	6–14	11.65	1.91	7–15

Note: Parental Education: 1 = “high school diploma or lower,” 2 = “some college,” 3 = “four year degree,” 4 = “graduate education and above.” Parental Income: 1 = ≤\$20K, 2 = \$20–\$40K, 3 = \$40–\$60K, 4 = \$60K+. CDI-C, Child Depression Inventory Self-Report; CEMS, Children’s Emotion Management Scale.

**p* < .05 significant chi-squared test of differences in ethnicity distribution between the 48 children who did complete the follow-up and the 12 who did not.

children with high PFC scores, children with no endorsed symptoms were recruited to establish a healthy comparison group (of interest in the current analyses). Approximately 6,000 checklists were distributed to sites between May 2003 and March 2005. In daycares and preschools, from which approximately 75% of the sample was ascertained, checklists were handed out to all parents of children in the target age range. Checklists were made available in waiting areas of primary care settings next to a poster describing a study of early emotional development. Completed checklists were collected by the sites and returned. Using this method, *N* = 1,474 checklists were returned, and those with scores of 0 (presumed healthy) or 3 or higher (above established cutoff) were contacted for further participation. Of those who returned checklists, *N* = 335 were ineligible due to being out of the age range and *N* = 240 had PFC scores out of range (e.g., 1 or 2). The remaining *N* = 899 met all initial screening and inclusion criteria and were contacted by phone for further screening. Based on phone screening, subjects with chronic illness, marked speech and language delays, and/or neurologic or autistic spectrum disorders were excluded. Those without exclusions (*N* = 416) were invited for study participation, and *N* = 305 agreed and presented for the assessment.

Psychiatrically healthy children and those with preschool-onset MDD who had no contraindication for MRI were eligible for an MRI scan session after the fourth wave of annual assessments. Of the 86 eligible and psychiatrically healthy children, 49 agreed to be scanned. Of these 49 children, 6 were excluded because they developed an Axis I psychiatric disorder. Of the remaining 43, data was excluded from 10 children because of incomplete scan runs or poor quality fMRI data. (Note that 77% usable functional data is not unusual for child studies, but it improved after the start of the

study as our methods for ensuring compliance and low movement improved.) Of these 33 children, 29 also had complete data on all of the clinical variables examined at scan.

The second wave of recruitment involved recruiting additional children with no prior psychiatric diagnosis to enhance the subsample of healthy subjects. Subjects were screened from local schools using the Child Behavior Checklist (Achenbach & Edelbrock, 1991), and those in the healthy range without scan exclusions were invited to participate. Forty-two children were interviewed using the same age-appropriate semistructured diagnostic measure used in the larger sample (see below for details). One child was excluded for a neurological issue and 4 were excluded because they met criteria for an Axis I psychiatric diagnosis. An additional 4 children were excluded because of poor-quality MRI data. Of the remaining 33 children, 31 had complete data on all of the clinical variables examined at scan.

Diagnostic assessment

Trained staff from the WUSM Early Emotional Development Program conducted up to three in-person assessment sessions with participants and their primary caregivers prior to the MRI scan session, and up to two more in-person assessments over the course of 3 years (to date) after the scan session. The assessments included age-appropriate, semistructured diagnostic interviews to assess psychiatric symptoms; before children were age 8, the Preschool-Age Psychiatric Assessment (Egger, Erkanli, Keeler, & Potts, 2006; Luby et al., 2009) was used, whereas parent and child report on the Childhood and Adolescent Psychiatric Assessment was used when children were older than age 8. Because the current report focuses on psychiatrically healthy children, any child who met criteria for any Axis I DSM-IV psychiatric disorder prior to the first scan session was excluded from the analyses.

Self-report instruments

Children completed the Children’s Depression Inventory (CDI-C; Kovacs, 1985) at the time of first scan to assess current depression symptom severity. Children then completed the CDI-C again at a follow-up approximately 18 months later. Total CDI-C scores were computed and converted to *t* scores. The Children’s Emotion Management Scale (CEMS) was used to measure children’s ability to manage/regulate their experiences of sadness (Zeman, Shipman, & Penza-Clyve, 2001). The CEMS was administered at the annual assessment prior to the scanning session and then again at the 18-month follow-up. For the CEMS, we focused on the sadness coping subscale as a relevant measure of emotion regulation skills, where higher CEMS sadness coping scores indicate better emotion regulation skills.

Functional task and stimuli

The facial emotion processing task of interest here represents one segment of a longer scanning session that the children in

the PDS completed, which included high-resolution structural imaging, diffusion imaging, and resting-state functional connectivity scans. The fMRI task was an event-related facial emotion processing task, similar to that used in previous research in adults and adolescents with depression (Beesdo et al., 2009; Gotlib et al., 2005; Lau et al., 2009; Roberson-Nay et al., 2006; Surguladze et al., 2005; Thomas et al., 2001). Children were shown faces that varied in affective content and were asked to decide whether the face was male or female. We used a task that did not require explicit attention to the emotional content because evidence suggests that heightened amygdala responses associated with depression may be more apparent with a less constrained response (Fales et al., 2008; Monk et al., 2008). The face stimuli were drawn from the MacArthur Network Face Stimuli Set (Tottenham et al., 2009). Children were shown sad, fearful, angry, happy, and neutral expressions from 10 individuals. In addition, we created intermediate sad, fearful, angry, and happy expressions by morphing the neutral expression for each individual with each of their emotional expressions so that the resulting face was halfway between neutral and the target emotion (Morph Age software). We included these stimuli to test whether brain activation biases in depression may be more apparent with lower intensity emotional expressions. Thus, each “actor” in the stimulus set provided a total of 9 expressions (neutral; and 50% and 100% sad, fearful, angry, and happy).

In this report, we focus on brain response to sad faces versus neutral faces, given prior findings suggesting that there are specific impairments associated with sad face processing in children with a history of preschool-onset depression from both this study sample as well as from an independent sample (Barch et al., 2012; Gaffrey, Barch, Singer, Shenoy, & Luby, 2013; Gaffrey et al., 2011). Further, we focused on the average functional activations to the full and half intensity faces in order to increase power and because our prior work in this sample suggests that alterations associated with depression are not specific to either intensity of sad expression (Barch et al., 2012). The results were not substantively different if we focused only on the high-intensity faces.

Each scan run consisted of 45 stimuli, 5 from each of the 9 conditions. Each stimulus was presented for 2500 ms, followed by an intertrial interval ranging between 500 and 6500 ms. Each child performed two task runs with no stimuli repetition. In addition, prior to performing this task, all children underwent a mood induction in the scanner based on prior work (Gotlib et al., 2005). We included this mood manipulation because affective processing biases (Scher, Ingram & Segal, 2005) and hyperactivity of limbic regions (Ramel et al., 2007) can be reactivated in individuals with a past history of MDD following a mood induction and can be elicited using mood induction in nondepressed children at risk due to maternal MDD history (Joormann & Gotlib, 2006; Joormann, Talbot, & Gotlib, 2007; Taylor & Ingram, 1999). We used a film clip from *My Girl* (see Joormann et al., 2007), which focuses on a child’s loss of a close friend, to induce a negative mood state, coupled with directions to elaborate on this negative mood by imagining oneself in the situation (Wester-

mann, Spies, Stahl, & Hesse, 1996). This mood induction occurred immediately before children began the facial emotion processing task. A positive mood repair clip was shown at the end of the session.

MRI data acquisition

Structural and functional scanning was performed on a 3.0 Tesla TIM TRIO Siemens whole-body system. For structural data, three-dimensional T1-weighted images (resonance time = 2400 ms, echo time = 3.16 ms, flip angle = 8°, slab = 176 mm, 176 slices, matrix size = 256 × 256, field of view = 256 mm, voxel size = 1 × 1 × 1 mm, 1 signal average) were acquired in the sagittal plane with the use of magnetization-prepared rapid gradient echo sequence. Blood oxygen level dependent (BOLD) images during face processing were acquired with a T2*-weighted asymmetric spin-echo echo-planar sequence (resonance time = 3000 ms, echo time = 27 ms, flip angle = 90°, field of view = 256 mm, voxel size = 4 × 4 × 4 mm) in the axial plane paralleling to the anterior–posterior commissure, with a 12-channel head coil. During each functional run, 99 sets of 36 contiguous axial images with isotropic voxels (4 mm³) were acquired parallel to the anterior–posterior commissure plane.

fMRI data processing

The fMRI data were preprocessed using standard preprocessing steps, including the following:

1. compensation for slice-dependent time shifts;
2. removal of the first five images of each run to allow BOLD signal to reach steady state;
3. elimination of odd/even slice intensity differences due to interpolated acquisition;
4. realignment of data acquired in each subject within and across runs to compensate for rigid body motion (Ojemann et al., 1997);
5. intensity normalization to a whole-brain mode value of 1,000;
6. registration of the three-dimensional structural volume (T1) to the atlas representative template in the Talairach coordinate system (Talairach & Tournoux, 1988) using a 12-parameter affine transform and resampling to 1 mm cubic representation (Buckner et al., 2004; Ojemann et al., 1997);
7. coregistration of the three-dimensional fMRI volume to the T2, and the T2 to the participants structural image;
8. transformation of the fMRI volumes to atlas space using a single affine 12-parameter transform; and
9. spatial smoothing using a 6-mm full-width at half-maximum Gaussian filter.

The common atlas template was optimized for children in our age range; the use of co-registration to this type of common template space has been validated in several prior studies (Burgund et al., 2002; Kang, Burgund, Lugar, Petersen, & Schlaggar, 2003).

Previously validated “motion scrubbing” procedures adapted for fMRI data were used to remove artifacts due to head motion (Power, Barnes, Snyder, Schlaggar, & Petersen, 2013; Power, Mitra, et al., 2013). As discussed and validated previously for use in this sample (Pagliaccio et al., 2013), the motion-scrubbing procedure assesses framewise displacement based on the parameters used in reprocessing Step 4. This represents the differential head motion from one acquisition frame to the next summing across linear (x , y , and z) and rotational displacements (yaw, pitch, and roll, where degrees of rotation are converted to millimeters of movement by calculating displacement on the surface of a sphere with a radius of 50 mm). Any frame with a sum displacement greater than 0.9 mm was masked out of the analysis.

Analysis of fMRI data was performed using in-house software (FIDL analysis package, <http://www.nil.wustl.edu/~fidl>) utilized in numerous previously published studies (Becerril, Repovs, & Barch, 2011; Braver, Paxton, Locke, & Barch, 2009; Gaffrey et al., 2011; Jimura, Locke, & Braver, 2010; Ollinger, Shulman, & Corbetta, 2001). Estimates of BOLD response to each face type for each subject were obtained using fixed effects general linear models incorporating regressors for linear trend and baseline shifts. A hemodynamic response shape was assumed (SPM canonical function) and used to derive magnitude estimates relative to fixation baseline. These single-subject estimates were then entered into group-level analyses that treated subjects as random effects.

A priori region of interest definitions

As a measure of responsivity to emotionally evocative stimuli, we examined responses to sad faces versus neutral faces in regions associated with emotion responsivity and in regions associated with emotion regulation. To do so, we examined task-related responsivity in two previously identified networks (Anticevic & Repovs, 2011; Luking et al., 2011). The first was a network of regions associated with bottom-up emotion responsivity, including the bilateral amygdala and regions that show positive resting-state functional connectivity with the amygdala (i.e., hippocampal, parahippocampal, striatal, and temporal regions). Although an oversimplification, we will refer to this as the emotion reactivity network (Table 2). The second network included regions associated with top-down emotion regulation that show negative resting-state functional connectivity with the amygdala, including the dorsal frontal and parietal regions. Again, though an oversimplification, we will refer to this as the emotion regulation network (Table 2). In prior work, we found that connectivity of both sets of regions was disturbed in children with a history of depression and in those at risk for depression (Luking et al., 2011) and are thus of key interest for studying responses to emotional faces. For each region, we created a spherical region of interest, 12 mm in diameter, around its centroid, listed in Table 2. Magnitude estimates of the response to sad versus neutral faces were averaged across

Table 2. Regions included as part of emotion reactivity and emotion regulation networks

Region	X	Y	Z
Emotion Reactivity Network			
Left			
Amygdala	-23	-8	-16
Hippocampus	-27	-26	-12
Inferior temporal gyrus	-41	-9	-29
Superior temporal gyrus	-27	6	-32
Right			
Amygdala	18	-8	-20
Hippocampus	34	-26	-14
Parahippocampal gyrus	18	-23	-14
Putamen	25	2	-7
Superior temporal gyrus	33	10	-28
Emotion Regulation Network			
Left			
Inferior parietal lobule (BA39)	-36	-61	40
Middle frontal gyrus (ant. BA9)	-37	28	32
Middle frontal gyrus (post. BA6)	-26	9	51
Precentral gyrus (BA6)	-43	-2	27
Superior frontal gyrus (ant. BA10)	-30	49	14
Superior frontal gyrus (post. BA8)	-4	15	49
Supramarginal gyrus	-53	-50	37
Right			
Inferior parietal lobule (BA40)	44	-42	53
Middle frontal gyrus (BA9)	32	31	30
Postcentral gyrus (BA2)	53	-27	42
Precentral gyrus (BA6)	51	-6	32
Precuneus (BA7)	-11	-73	42

Note: BA, Brodmann area.

all of the regions of interest within each of the two networks to create an overall estimate of responses in emotion reactivity regions and in emotion regulation regions.

Structural data processing

We were interested in the volume of the amygdala, hippocampus, and insula given evidence for altered volume of these regions in depression and given their putative involvement in emotion processing. FreeSurfer version 4.5.0 (<http://surfer.nmr.mgh.harvard.edu>; Fischl et al., 2002, 2004) was used to segment each participant’s anatomical image (using the Desikan-Killiany Atlas for cortical segmentation; Desikan et al., 2006), allowing estimation of the volume of the left and right hippocampal, amygdala, and insula. To reduce the number of predictors in our analyses, we averaged volume estimates for both hemispheres of the amygdala, hippocampus, and insula. In addition, we estimated whole-brain volume (total gray + cortical white matter volume). The white and pial FreeSurfer surfaces were visually inspected and were regenerated with manual intervention to correct errors when necessary. Cortical gray matter volume was defined as the volume between the pial and white matter surfaces.

White matter volume was calculated by subtracting the sub-cortical and ventricular volumes from the volume bounded by the white matter surface.

Analytical approach

The aim of the current analyses was to investigate how the structure and function of key brain regions involved in emotion reactivity/regulation predict concurrent and follow-up measures of subclinical depressive symptomology and emotion regulation skills in psychiatrically healthy children. First, we used linear regression models to test whether amygdala volume, hippocampal volume, insula volume, emotion reactivity network activity, and/or emotion regulation network activity predict individual differences in self-reported depression (CDI-C) or self-reported emotion regulation skills (CEMS) from the assessments concurrent with the imaging data acquisition. To test the specificity of the brain volume effects, we controlled for effects of whole-brain volume as well. Second, we used linear regressions to test whether the factors that significantly predicted scores at scan also predict CDI-C or CEMS scores at the 18-month follow-up. When significant predictors were found, we then tested whether these factors remained significant when controlling for CDI-C or CEMS scores concurrent to the scan (i.e., to test for factors that predict change over time). Standardized regression coefficients and their associated *T* statistics and *p* values are presented in Table 3 and in the text below.

Results

Demographic and clinical characteristics

Comparing the children recruited as part of the two recruitment waves, there were no significant differences in age, $t(58) = 0.35, p = .58$; sex, $\chi^2(1) = 0.61, p = .38$; ethnicity, $\chi^2(2) = 2.65, p = .27$; CDI-C at scan, $t(58) = -0.77, p = .44$; activity in the emotion regulation network, $t(58) = -0.31, p = .76$; or any brain volume measures of interest here ($ps > .05$). Children recruited in the first wave did show somewhat higher CEMS scores at scan, $t(58) = 2.28, p = .03$, and somewhat lower activity in the emotion reactivity network, $t(58) = -2.50, p = .02$.

Table 1 summarizes the demographic and self-report information from the children who contributed data at the time of scan ($N = 60$) and from the children who contributed data to the 18-month follow-up ($N = 48$). There were no significant differences in sex, $\chi^2(1) = 0.70, p = .47$; age, $t(58) = 0.67, p = .51$; family income, $t(58) = 1.71, p = .09$; parental education level, $t(58) = 1.63, p = .11$; recruitment wave, $\chi^2(1) = 0.60, p = .44$; CDI-C scores, $t(58) = -0.23, p = .82$; CEMS sadness coping scores, $t(58) = 0.88, p = .38$; or any of the brain structure/function measures ($ps > .05$) between the 48 children who did and the 12 children who did not complete the follow-up. None of these factors predicted the likelihood of attrition in a logistic regres-

Table 3. Regressions predicting CDI-C and CEMS self-reports concurrent to scan

	β	<i>T</i>	<i>p</i>
Dependent Measure: CDI-C Child Report $F(6, 53) = 4.30, p = .001, \text{Adj. } R^2 = .25$			
Bilateral			
Amygdala volume	-0.09	-0.64	0.53
Hippocampal volume	-0.31	-2.01	0.04
Insula volume	0.31	1.93	0.06
BOLD response to sad vs. neutral faces in			
Emotion reactivity network	0.41	3.40	0.001
Emotion regulation network	-0.02	-0.17	0.86
Whole brain volume	-0.16	0.90	0.37
Dependent Measure: CEMS Sadness Coping $F(6, 53) = 3.85, p = .003, \text{Adj. } R^2 = .23$			
Bilateral			
Amygdala volume	-0.36	-2.50	0.02
Hippocampal volume	0.19	1.22	0.23
Insula volume	0.37	2.30	0.03
BOLD response to sad vs. neutral faces in			
Emotion reactivity network	-0.28	-2.23	0.03
Emotion regulation network	0.10	0.81	0.42
Whole brain volume	0.08	0.45	0.65

Note: CDI-C, Child Depression Inventory Self-Report; CEMS, Children's Emotion Management Scale; BOLD, blood oxygen level dependent.

sion ($ps > .05$). However, the follow-up included a smaller percentage of Caucasian children, $\chi^2(2) = 7.86, p = .02$, because 11 of the 12 children who did not complete the follow-up were Caucasian.

Intercorrelation among predictors and network reliability

Examining the correlation between our two outcome measures of interest (see Table 4), we found that CDI-C and CEMS scores were not significantly correlated at the time of scan, $r(58) = -.19, p = .146$, or at follow-up, $r(46) = -.16, p = .28$. However, as expected, CDI-C scores were significantly correlated across time from scan to follow-up, $r(46) = .55, p < .001$, as were CEMS scores, $r(46) = .38, p = .008$.

Examining our predictors of interest (see Table 4), we found significant intercorrelation among insula, hippocampal, and amygdala volumes, insula-hippocampus: $r(58) = .55, p < .001$; insula-amygdala: $r(58) = .43, p = .001$; hippocampus-amygdala: $r(58) = .48, p < .001$. We also found that our measures of average network activity were highly reliable because there was strong intercorrelation among activity estimates from the emotion reactivity network regions of interest (Cronbach $\alpha = 0.77$) and among emotion regulation network regions of interest (Cronbach $\alpha = 0.90$). In addition, average sad versus fearful face activity estimates from the two networks were correlated, $r(58) = .35, p = .006$. There were

Table 4. Intercorrelations between factors of interest

	CDI-C Follow-Up	CEMS Scan	CEMS Follow-Up	Insula	Hippocampus	Amygdala	Emotion Reactivity Network	Emotion Regulation Network
CDI-C								
Scores at scan	0.55**	-0.19	-0.32*	-0.02	-0.33**	-0.23	0.44**	0.18
Scores at follow-up (<i>N</i> = 48)		-0.36*	-0.16	-0.18	-0.33*	-0.17	0.02	-0.17
CEMS								
Scores at scan			0.38**	0.38**	0.29*	-0.06	-0.23	0.06
Scores at follow-up (<i>N</i> = 48)				0.38**	0.46**	0.21	-0.31*	-0.13
Bilateral								
Insula volume					0.55**	0.43**	0.00	0.04
Hippocampal volume						0.48**	-0.10	-0.09
Amygdala volume							-0.06	-0.16
Emotion reactivity network activity								0.35**

* $p < .05$. ** $p < .01$.

no significant correlations between insula, hippocampal, or amygdala volumes and activity in either network ($ps > .05$).

Predicting concurrent self-reports of subclinical depressive symptomology and emotion regulation

CDI-C scores. A linear regression with amygdala volume, hippocampus volume, insula volume, and activity in the emotion reactivity and emotion regulation networks predicted a significant amount of variance in self-reported subclinical depressive symptomology (CDI-C scores) at the time of scan, $F(6, 53) = 4.30$, $p = .001$, adjusted $R^2 = .25$. As shown in Table 3, both reduced hippocampus volume and increased BOLD response to sad versus neutral faces in the emotion reactivity network predicted higher self-reports of depressive symptoms, though activity in the emotion regulation network did not predict it. It is important that both hippocampus volume and the emotion reactivity network activity significantly predicted CDI-C scores, while controlling for whole-brain volume, which was not a significant predictor.

CEMS scores. As above, we used a linear regression to test whether brain volume and/or reactivity measures predict concurrent self-reports of emotion regulation on the CEMS (higher scores on the CEMS indicating better sadness coping skills). This regression predicted a significant amount of variance in CEMS scores, $F(6, 53) = 3.85$, $p = .003$, adjusted $R^2 = .23$. As shown in Table 3, larger insula volume and smaller amygdala volume predicted better emotion regulation skills. In addition, less BOLD response to sad versus neutral faces in the emotion reactivity network predicted higher CEMS scores, though activity in the emotion regulation network did not. It is important that these measures significantly predicted CEMS scores, controlling for whole-brain volume, which was not a significant predictor.

Predicting follow-up self-reports of subclinical depressive symptomology and emotion regulation

CDI-C scores. We tested whether the same structural and functional measures that predicted concurrent CDI-C scores could also predict CDI-C scores at the 18-month follow-up. Smaller hippocampal volumes predicted higher CDI-C scores at follow-up ($\beta = -0.33$, $t = -2.34$, $p = .02$), but emotion reactivity network activity did not ($\beta = 0.02$, $t = 0.12$, $p = .91$). However, hippocampal volume did not predict CDI-C scores at follow-up when accounting for concurrent CDI scores ($\beta = -0.15$, $t = -1.11$, $p = .27$).

CEMS scores. As above, we tested whether the same structural and functional measures that predicted concurrent CEMS scores also predicted CEMS scores at follow-up. Larger insula volumes ($\beta = 0.38$, $t = 2.79$, $p = .008$) and less BOLD response to sad versus neutral faces in the emotion reactivity network ($\beta = -0.311$, $t = -2.22$, $p = .031$) both predicted better CEMS sadness coping scores at follow-up. However, amygdala volumes did not ($\beta = 0.21$, $t = 1.45$, $p = .15$). It is important that, as shown in Figure 1, insula volumes continued to predict CEMS scores at follow-up even after accounting for concurrent CEMS scores ($\beta = 0.29$, $t = 2.11$, $p = .04$), suggesting that greater insula volumes predict a greater increase in self-reported emotion regulation over time. There was also a trend for less BOLD activity in the emotion reactivity network to predict CEMS scores at follow-up even after accounting for concurrent CEMS scores ($\beta = -0.23$, $t = -1.69$, $p = .098$).

Discussion

The current findings demonstrate specific brain-behavioral relationships in emotion processing relevant to depressive




Figure 1. Scatterplot illustrating the relationship between insula volumes and emotion regulation at 18-month follow-up. The Children's Emotional Management Scale sadness coping values represent the standardized residuals of follow-up score accounting for baseline Children's Emotional Management Scale sadness coping scores, where positive scores indicate improvements in sadness coping from baseline to follow-up. This relationship remains significant ($\beta = 0.29$, $t = 2.10$, $p = .04$) after Winsorizing the volume of the potential outlier participant with the largest insula volume (11852.5 mm³).

symptoms in the normative subclinical range and emotion regulation in healthy children. We found evidence that brain structure (decreased hippocampal volume) and function (increased sad-neutral face activity in the emotion reactivity network) relate to increased subclinical depressive symptomatology in healthy school-age children at the time of scan, while increased amygdala volume, increased emotion reactivity network activity, and decreased insula volume relate to better emotion regulation skills. Furthermore, brain structure (increased insula volume) and function (decreased emotion reactivity network activity) can prospectively predict improvements in emotional regulation skills (CEMS sadness coping scores) at an 18-month follow-up. The volume of the insula was predictive of later regulation skills over and above the effect of emotion regulation scores at scan.

These findings begin to elucidate how the structure of the several key brain regions, the hippocampus, amygdala, and insula, and the overall responsiveness of emotion reactivity regions are associated with the normative experience of depressive symptoms and with emotion regulatory capacities as early as school age. This type of multimodal individual difference approach in a psychiatrically healthy, developmental sample can be highly informative for studies of pediatric and adult psychopathology. Such investigations in healthy developing populations help to inform our understandings of brain-behavior relationships underlying normal variation in emotionality. Such data may guide our understanding of the normative developmental trajectories of these relationships. Furthermore, understanding normal relationships between brain

structure/function and emotion can then lead to more informed interpretations of neural alterations in depression; that is, if increased insula volume positively predicts sadness coping skills and improvements in coping skills over development, then decreased insula volumes evident in depressed patients versus healthy controls may indicate emotion regulation deficits in the depressed sample.

Predicting subclinical depressive symptomatology

Our finding that decreased hippocampal volumes predict greater self-reported depressive symptomatology among healthy children is of interest because reports of decreased volume in pediatric depression are somewhat mixed. This finding extends the literature by suggesting that the severity of depressive symptomatology, even those not reaching clinically significant thresholds, is related to reductions in hippocampal volume in childhood. Although hippocampal volume was predictive of depressive symptomatology at ~18 months postscan, this did not predict over and above CDI-C scores concurrent to the scan. Thus, while hippocampal volume may be a key marker of or risk factor for depressive symptomatology, volume does not appear to relate significantly to change in depressive symptomatology over this relatively short interval. As such, this finding might suggest that hippocampal volume reductions index some early occurring vulnerability factor that may not change over time. However, an important follow-up to this is to explore whether change in hippocampal volume over time relates to change in depressive symptomatology over longer periods of time.

Sad versus neutral face activity in emotion reactivity regions was a robust predictor of depressive symptomatology, where greater response magnitude predicted greater depressive symptomatology. This is highly congruous with studies suggesting that hyperreactivity to negative emotional stimuli in the amygdala, hippocampus, and related regions is associated with depression in both adult and pediatric populations. This finding elaborates on previous work to suggest that individual differences in severity may be reflected in reactivity levels across this whole network of regions that show positive resting-state functional connectivity with the amygdala. As noted previously, this effect in psychiatrically healthy children does indicate that this may be a normative brain-behavior relationship, not restricted to clinical populations. Of interest here, the magnitude of activity in this network did not predict follow-up depression severity. Thus, activity in emotion reactivity regions may reflect more state-level differences in depressive symptomatology rather than trait-level differences or trajectories over time. Relevant to this, research on the test-retest reliability of amygdala reactivity to emotional faces has shown relatively low values (over a 90-day gap, moderate reliability for fearful faces: ICC ~ 0.3 to 0.4; low reliability for angry faces: ICC ~ -0.1 to 0.1; Sauder, Hajcak, Angstadt, & Phan, 2013). This could reflect changes in amygdala reactivity as a function of emotional state, and if so, it may suggest that these reactivity measures may be better state-level predictors.

Predicting emotion regulation skills

Turning to emotion regulation (CEMS scores of sadness coping skills), we found that larger amygdala volumes predicted poorer emotion regulation skills at the time of scan. Though literature on adult depression shows mixed results for amygdala volume effects, the pediatric literature, structure–function work, and animal studies suggest that larger amygdala volumes are related to both early adversity and greater emotionality and negative outcomes (e.g., Tottenham et al., 2010; van der Plas et al., 2010). For example, girls, particularly those with a family history of depression, show a positive correlation between amygdala volume and fearfulness (van der Plas et al., 2010). Our results build on these and other results to suggest that larger amygdala volumes are related to less effective self-reported sadness coping skills, even in healthy school-age children.

Greater activity in emotion reactivity network regions was related to poorer concurrent emotion regulation skills, though it only predicted change in emotion regulation over time at a trend level. Previous work in this sample also showed that greater right amygdala sad versus neutral face activity in school-age children was related to poorer sadness regulation skills summing across all three subscales of the CEMS (Pagliaccio et al., 2013). This is highly relevant for the well-replicated finding that people with depression show elevated amygdala responses to emotional stimuli and that greater depression severity correlates with greater amygdala responses to these stimuli (e.g., Barch et al., 2012; Beesdo et al., 2009; Gaffrey et al., 2011; Pagliaccio et al., 2011; Yang et al., 2010). First, these findings suggest that exploring individual differences in constructs, like emotion regulation, may be key to understanding the underlying neural and psychological deficits in depression (Aldao, Nolen-Hoeksema, & Schweizer, 2010; Garnefski & Kraaij, 2006; Joormann & Gotlib, 2010; Silk, Steinberg, & Morris, 2003). Second, this finding indicates that exploring activity in a variety of emotion reactivity-related regions may help to give a more holistic understanding of the neural correlates of emotional functioning rather than focusing, for example, solely on the amygdala.

Finally, we found that larger insula volumes predicted better emotion regulation skills at scan and improvements in emotion regulation over time (i.e., at 18-month follow-up, controlling for scan concurrent scores). This is a key finding given work highlighting decreased insula volume as a marker of depression in adults (Lai & Wu, 2014; Liu et al., 2014; Peng et al., 2011) and in children (Belden et al., 2014). Further, while much of the literature has focused on the insula's relationship with guilt, disgust, and somatosensory experience, these findings suggest that insula volume may also show a relationship with emotion regulation skills. Consistent with this, pathological guilt may be viewed as a deficit of emotion regulation of sadness (Cole & Michel, 1994). The finding that insula volume can prospectively predict improvement in emotion regulation is particularly important to improving our understandings of normative developmental trajectories of emotionality and also may be key to

identifying neural markers of negative trajectories contributing to risk for developing depression.

Limitations

One limitation to this study is the sample size. Though our sample size is relatively large for neuroimaging studies, greater sample sizes are integral to exploring individual differences relationships for both cross-sectional and longitudinal analyses. In contrast, the current subsample of children from the PDS have been thoroughly screened and evaluated in great detail, which helps to establish, through several waves of clinical interview, that our participants are truly psychiatrically healthy. In addition, there are likely other important brain–behavior relationships not being captured by the current analyses, but we aimed to maintain a very specific focus on brain regions well characterized to relate to emotion reactivity/regulation and to show disruptions in MDD. Next, while the prospective longitudinal design utilized here is quite powerful for exploring developmental change, the relatively high correlation between CDI-C or CEMS scores concurrent to the scan and at later follow-up may limit our ability to find predictors that account for variance in follow-up scores over and above the predictive power of scores concurrent with the scan data. Given the null effects of emotion regulation network activity and the positive correlation between activity in both networks, another limitation to note is that only activity during an emotional face-processing task was utilized here. While these regions are likely required even for implicit regulation of induced emotion, using a task requiring more explicit regulation may tap into key individual differences in emotion regulation-related brain activity and may reveal stronger relationships to depression and emotion regulation either concurrently or over time.

A further conceptual and methodological limitation to this work and to most studies relating neuroimaging to emotion and psychopathology is that they often require reducing the many complexities of emotion (e.g., tone, intensity, time course, regulation) into simplified summary statistics, such as questionnaire scores, for the ease of statistical analysis. In parallel, we must generally reduce the complexity of the dynamic and developing neural circuitry underlying emotion into summary measures of volume or reactivity, which again are useful for statistical analysis. Thus, while the current approaches are still powerful for understanding brain–behavior relationships, they are potentially limited by the current measures, which may be masking key nuances in phenotypic variation and internal emotional experience. Hopefully, future work can continue to expand our understandings of these relationships and shed light on the developmental psychopathology of mood disorders by exploring the great complexity of both emotional experience and its underlying neurobiology.

Future directions

In order to more fully understand the neural mechanisms that underlie individual differences in emotion reactivity/regula-

tion, work must continue to explore brain–behavior relationships, not only those relationships that may be disrupted in psychopathology, but also relationships during normative development. This requires more study of individual differences in processes/skills that are altered in MDD and other disorders in relation to brain structure and function, as well as in relation to risk/resilience factors. In addition, it will be important to test what neural markers help to explain further longitudinal change over time, exploring more factors beyond depressive

symptomology and sadness coping, and prospectively beyond this 18-month follow-up. Furthermore, it will be important to explore how *trajectories* of neural change predict emotional development. This will hopefully be addressed by data from the PDS sample and others in the future. In addition, it will be important to examine behavioral and neural markers of risk for depression even earlier in development, as the tools and methods for noninvasive brain imaging in young children become more robust and easy to implement.

References

- Achenbach, T. M., & Edelbrock, C. (1991). *Manual for the Child Behavior Checklist—Teacher's Report Form*. Burlington, VT: University of Vermont, Department of Psychiatry.
- Aldao, A., Nolen-Hoeksema, S., & Schweizer, S. (2010). Emotion-regulation strategies across psychopathology: A meta-analytic review. *Clinical Psychology Review, 30*, 217–237.
- Amico, F., Meisenzahl, E., Koutsouleris, N., Reiser, M., Möller, H.-J., & Frodl, T. (2011). Structural MRI correlates for vulnerability and resilience to major depressive disorder. *Journal of Psychiatry and Neuroscience, 36*, 15–22.
- Andersen, S. L., & Teicher, M. H. (2004). Delayed effects of early stress on hippocampal development. *Neuropsychopharmacology, 29*, 1988–1993.
- Anticevic, A., & Repovs, G. (2011). Emotion effects on attention, amygdala activation, and functional connectivity in schizophrenia. *Schizophrenia Bulletin, 38*, 967–980.
- Anticevic, A., Repovs, G., & Barch, D. M. (2010). Resisting emotional interference: Brain regions facilitating working memory performance during negative distraction. *Cognitive, Affective, & Behavioral Neuroscience, 10*, 159–173.
- Barch, D. M., Gaffrey, M. S., Botteron, K. N., Belden, A. C., & Luby, J. L. (2012). Functional brain activation to emotionally valenced faces in school-aged children with a history of preschool-onset major depression. *Biological Psychiatry, 72*, 1035–1042.
- Beesdo, K., Lau, J., Guyer, A., McClure-Tone, E., Monk, C., Nelson, E., et al. (2009). Common and distinct amygdala-function perturbations in depressed vs. anxious adolescents. *Archives of General Psychiatry, 66*, 275.
- Belden, A., Barch, D. M., Oakberg, T. J., April, L. M., Harms, M. P., Botteron, K. N., et al. (2014). *Anterior insula volume and guilt: Neurobehavioral markers of recurrence after early childhood major depressive disorder*. Manuscript submitted for publication.
- Belden, A. C., Luby, J. L., Pagliaccio, D., & Barch, D. M. (2014). Neural activation associated with the cognitive emotion regulation of sadness in healthy children. *Developmental Cognitive Neuroscience, 9C*, 136–147.
- Becerril, K. E., Repovs, G., & Barch, D. M. (2011). Error processing network dynamics in schizophrenia. *NeuroImage, 54*, 1495–1505.
- Birmaher, B., Ryan, N. D., Williamson, D. E., & Brent, D. A. (1996). Childhood and adolescent Depression: A review of the past 10 years: Part I. *Journal of the American Academy of Child & Adolescent Psychiatry, 35*, 1427–1439.
- Braver, T., Paxton, J., Locke, H., & Barch, D. (2009). Flexible neural mechanisms of cognitive control within human prefrontal cortex. *Proceedings of the National Academy of Sciences, 106*, 7351.
- Buckner, R. L., Head, D., Parker, J., Fotenos, A. F., Marcus, D., Morris, J. C., et al. (2004). A unified approach for morphometric and functional data analysis in young, old, and demented adults using automated atlas-based head size normalization: Reliability and validation against manual measurement of total intracranial volume. *NeuroImage, 23*, 724–738.
- Buhle, J. T., Silvers, J. A., Wager, T. D., Lopez, R., Onyemekwu, C., Kober, H., et al. (2013). Cognitive reappraisal of emotion: A meta-analysis of human neuroimaging studies. *Cerebral Cortex*. Advance online publication.
- Burgund, E. D., Kang, H. C., Kelly, J. E., Buckner, R. L., Snyder, A. Z., Petersen, S. E., et al. (2002). The feasibility of a common stereotactic space for children and adults in fMRI studies of development. *NeuroImage, 17*, 184–200.
- Burke, H. M., Davis, M. C., Otte, C., & Mohr, D. C. (2005). Depression and cortisol responses to psychological stress: A meta-analysis. *Psychoneuroendocrinology, 30*, 846–856.
- Burke, K. C., Burke, J. D., Rae, D. S., & Regier, D. A. (1991). Comparing age at onset of major depression and other psychiatric disorders by birth cohorts in five US community populations. *Archives of General Psychiatry, 48*, 789–795.
- Caetano, S. C., Fonseca, M., Hatch, J. P., Olvera, R. L., Nicoletti, M., Hunter, K., et al. (2007). Medial temporal lobe abnormalities in pediatric unipolar depression. *Neuroscience Letters, 427*, 142–147.
- Campbell, S., & MacQueen, G. (2004). The role of the hippocampus in the pathophysiology of major depression. *Journal of Psychiatry and Neuroscience, 29*, 417–426.
- Canli, T., Cooney, R., Goldin, P., Shah, M., Sivers, H., Thomason, M., et al. (2005). Amygdala reactivity to emotional faces predicts improvement in major depression. *NeuroReport, 16*, 1267.
- Carballedo, A., Lisiecka, D., Fagan, A., Saleh, K., Ferguson, Y., Connolly, G., et al. (2012). Early life adversity is associated with brain changes in subjects at family risk for depression. *World Journal of Biological Psychiatry, 13*, 569–578.
- Chen, M. C., Hamilton, J. P., & Gotlib, I. H. (2010). Decreased hippocampal volume in healthy girls at risk of depression. *Archives of General Psychiatry, 67*, 270–276.
- Cole, P., & Michel, M. (1994). The development of emotion regulation and dysregulation: A clinical perspective. *Monographs of the Society for Research in Child Development, 59*, 73–100.
- Connolly, C. G., Wu, J., Ho, T. C., Hoefl, F., Wolkowitz, O., Eisendrath, S., et al. (2013). Resting-state functional connectivity of subgenual anterior cingulate cortex in depressed adolescents. *Biological Psychiatry, 74*, 898–907.
- Dannlowski, U., Kugel, H., Huber, F., Stuhrmann, A., Redlich, R., Grotegerd, D., et al. (2012). Childhood maltreatment is associated with an automatic negative emotion processing bias in the amygdala. *Human Brain Mapping, 11*, 2899–2909.
- Davidson, R., & Irwin, W. (1999). The functional neuroanatomy of emotion and affective style. *Trends in Cognitive Sciences, 3*, 11–21.
- Desikan, R. S., Ségonne, F., Fischl, B., Quinn, B. T., Dickerson, B. C., Blacker, D., et al. (2006). An automated labeling system for subdividing the human cerebral cortex on MRI scans into gyral based regions of interest. *NeuroImage, 31*, 968–980.
- Diekhof, E. K., Geier, K., Falkai, P., & Gruber, O. (2011). Fear is only as deep as the mind allows: A coordinate-based meta-analysis of neuroimaging studies on the regulation of negative affect. *NeuroImage, 58*, 275–285.
- Egger, H. L., Erkanli, A., Keeler, G., & Potts, E. (2006). Test–retest reliability of the Preschool Age Psychiatric Assessment (PAPA). *Journal of the American Academy of Child & Adolescent Psychiatry, 45*, 538–549.
- Fales, C., Barch, D., Rundle, M., Mintun, M., Snyder, A., Cohen, J., et al. (2008). Altered emotional interference processing in affective and cognitive-control brain circuitry in major depression. *Biological Psychiatry, 63*, 377–384.
- Fischl, B., Salat, D. H., Busa, E., Albert, M., Dieterich, M., Haselgrove, C., et al. (2002). Whole brain segmentation: Automated labeling of neuroanatomical structures in the human brain. *Neuron, 33*, 341–355.
- Fischl, B. B., Salat, D. H. D., van der Kouwe, A. J. W. A., Makris, N. N., Ségonne, F. F., Quinn, B. T. B., et al. (2004). Sequence-independent segmentation of magnetic resonance images. *NeuroImage, 23*(Suppl. 1), S69–S84.
- Fitzgerald, P. B., Laird, A. R., Maller, J., & Daskalakis, Z. J. (2008). A meta-analytic study of changes in brain activation in depression. *Human Brain Mapping, 29*, 683–695.
- Frodl, T., Jäger, M., Smajstrlova, I., Born, C., Bottlender, R., Palladino, T., et al. (2008). Effect of hippocampal and amygdala volumes on clinical

- outcomes in major depression: A 3-year prospective magnetic resonance imaging study. *Journal of Psychiatry and Neuroscience*, 33, 423–430.
- Fu, C. H. Y., Williams, S. C. R., Cleare, A. J., Brammer, M. J., Walsh, N. D., Kim, J., et al. (2004). Attenuation of the neural response to sad faces in major depression by antidepressant treatment: A prospective, event-related functional magnetic resonance imaging study. *Archives of General Psychiatry*, 61, 877–889.
- Fusar-Poli, P., Placentino, A., Carletti, F., Landi, P., Allen, P., Surguladze, S., et al. (2009). Functional atlas of emotional faces processing: A voxel-based meta-analysis of 105 functional magnetic resonance imaging studies. *Journal of Psychiatry and Neuroscience*, 34, 418–432.
- Gabard-Durnam, L. J., Flannery, J., Goff, B., Gee, D. G., Humphreys, K. L., Telzer, E., et al. (2014). The development of human amygdala functional connectivity at rest from 4 to 23 years: A cross-sectional study. *NeuroImage*, 95, 193–207.
- Gaffrey, M. S., Barch, D. M., Singer, J., Shenoy, R., & Luby, J. L. (2013). Disrupted amygdala reactivity in depressed 4- to 6-year-old children. *Journal of the American Academy of Child & Adolescent Psychiatry*, 52, 737–746.
- Gaffrey, M. S., Luby, J. L., Belden, A. C., Hirshberg, J. S., Volsch, J., & Barch, D. M. (2011). Association between depression severity and amygdala reactivity during sad face viewing in depressed preschoolers: An fMRI study. *Journal of Affective Disorders*, 129, 364–370.
- Gamefski, N., & Kraaij, V. (2006). Relationships between cognitive emotion regulation strategies and depressive symptoms: A comparative study of five specific samples. *Personality and Individual Differences*, 40, 1659–1669.
- Gotlib, I. H., Sivers, H., Gabrieli, J. D. E., Whitfield-Gabrieli, S., Goldin, P., Minor, K. L., et al. (2005). Subgenual anterior cingulate activation to valenced emotional stimuli in major depression. *NeuroReport*, 16, 1731.
- Green, J. G., McLaughlin, K. A., Berglund, P. A., Gruber, M. J., Sampson, N. A., Zaslavsky, A. M., et al. (2010). Childhood adversities and adult psychiatric disorders in the national comorbidity survey replication: I. Associations with first onset of DSM-IV disorders. *Archives of General Psychiatry*, 67, 113–123.
- Greicius, M., Flores, B., Menon, V., Glover, G., Solvason, H., Kenna, H., et al. (2007). Resting-state functional connectivity in major depression: Abnormally increased contributions from subgenual cingulate cortex and thalamus. *Biological Psychiatry*, 62, 429–437.
- Groenewold, N. A., Opmeer, E. M., de Jonge, P., Aleman, A., & Costafreda, S. G. (2012). Emotional valence modulates brain functional abnormalities in depression: Evidence from a meta-analysis of fMRI studies. *Neuroscience and Biobehavioral Reviews*, 37, 152–163.
- Gu, X., Hof, P. R., Friston, K. J., & Fan, J. (2013). Anterior insular cortex and emotional awareness. *Journal of Comparative Neurology*, 521, 3371–3388.
- Gur, R. C., Schroeder, L., Turner, T., McGrath, C., Chan, R. M., Turetsky, B. I., et al. (2002). Brain activation during facial emotion processing. *NeuroImage*, 16, 651–662.
- Guyer, A. E., Monk, C. S., McClure-Tone, E. B., Nelson, E. E., Roberson-Nay, R., Adler, A. D., et al. (2008). A developmental examination of amygdala response to facial expressions. *Journal of Cognitive Neuroscience*, 20, 1565–1582.
- Hamilton, J. P., Etkin, A., Furman, D. J., Lemus, M. G., Johnson, R. F., & Gotlib, I. H. (2012). Functional neuroimaging of major depressive disorder: A meta-analysis and new integration of baseline activation and neural response data. *American Journal of Psychiatry*, 169, 693–703.
- Hastings, R. S. R., Parsey, R. V. R., Oquendo, M. A. M., Arango, V. V., & Mann, J. J. (2004). Volumetric analysis of the prefrontal cortex, amygdala, and hippocampus in major depression. *Neuropsychopharmacology*, 29, 952–959.
- Heuser, I., Yassouridis, A., & Holsboer, F. (1994). The combined dexamethasone/CRH test: A refined laboratory test for psychiatric disorders. *Journal of Psychiatric Research*, 28, 341–356.
- Jacobson, L., & Sapolsky, R. (1991). The role of the hippocampus in feedback regulation of the hypothalamic–pituitary–adrenocortical axis. *Endocrine Reviews*, 12, 118–134.
- Jimura, K., Locke, H. S., & Braver, T. S. (2010). Prefrontal cortex mediation of cognitive enhancement in rewarding motivational contexts. *Proceedings of the National Academy of Sciences*, 107, 8871–8876.
- Joormann, J., & Gotlib, I. H. (2006). Is this happiness I see? Biases in the identification of emotional facial expressions in depression and social phobia. *Journal of Abnormal Psychology*, 115, 705–714.
- Joormann, J., & Gotlib, I. H. (2010). Emotion regulation in depression: Relation to cognitive inhibition. *Cognition and Emotion*, 24, 281–298.
- Joormann, J., Talbot, L., & Gotlib, I. H. (2007). Biased processing of emotional information in girls at risk for depression. *Journal of Abnormal Psychology*, 116, 135–143.
- Kang, H. C., Burgund, E. D., Lugar, H. M., Petersen, S. E., & Schlaggar, B. L. (2003). Comparison of functional activation foci in children and adults using a common stereotaxic space. *NeuroImage*, 19, 16–28.
- Kaufman, J., Martin, A., King, R., & Charney, D. (2001). Are child-, adolescent-, and adult-onset depression one and the same disorder? *Biological Psychiatry*, 49, 980–1001.
- Kessler, R., Berglund, P., & Demler, O. (2003). The epidemiology of major depressive disorder. *Journal of the American Medical Association*, 290, 1991–1992.
- Kessler, R. C., & Magee, W. J. (2009). Childhood adversities and adult depression: Basic patterns of association in a US national survey. *Psychological Medicine*, 23, 679.
- Koban, L., & Pourtois, G. (in press). Brain systems underlying the affective and social monitoring of actions: An integrative review. *Neuroscience & Biobehavioral Reviews*.
- Kovacs, M. (1985). The Children's Depression Inventory (CDI). *Psychopharmacology Bulletin*, 21, 995–998.
- Lai, C. H., & Wu, Y. T. (2014). Frontal-insula gray matter deficits in first-episode medication-naïve patients with major depressive disorder. *Journal of Affective Disorders*, 160, 74–79.
- Lamm, C., & Singer, T. (2010). The role of anterior insular cortex in social emotions. *Brain Structure and Function*, 214, 579–591.
- Lange, C., & Irle, E. (1999). Enlarged amygdala volume and reduced hippocampal volume in young women with major depression. *Psychological Medicine*, 34, 1059–1064.
- Lau, J. Y. F., Burt, M., Leibenluft, E., Pine, D. S., Rijdsdijk, F., Shiffrin, N., et al. (2009). Individual differences in children's facial expression recognition ability: The role of nature and nurture. *Developmental Neuropsychology*, 34, 37–51.
- Liu, C.-H., Jing, B., Ma, X., Xu, P.-F., Zhang, Y., Li, F., et al. (2014). Voxel-based morphometry study of the insular cortex in female patients with current and remitted depression. *Neuroscience*, 262, 190–199.
- Luby, J., Belden, A., Botteron, K., Marrus, N., Harms, M. P., Babb, C., et al. (2013). The effects of poverty on childhood brain development: The mediating effect of caregiving and stressful life events. *JAMA Pediatrics*, 167, 1135–1142.
- Luby, J. L., Heffelfinger, A., Koenig-McNaught, A. L., Brown, K., & Spitznagel, E. (2004). The Preschool Feelings Checklist: A brief and sensitive screening measure for depression in young children. *Journal of the American Academy of Child & Adolescent Psychiatry*, 43, 708–717.
- Luby, J. L., Si, X., Belden, A., Tandon, M., & Spitznagel, E. (2009). Preschool depression: Homotypic continuity and course over 24 months. *Archives of General Psychiatry*, 66, 897.
- Luking, K. R., Repovs, G., Belden, A. C., Gaffrey, M. S., Botteron, K. N., Luby, J. L., et al. (2011). Functional connectivity of the amygdala in early-childhood-onset depression. *Journal of the American Academy of Child & Adolescent Psychiatry*, 50, 1027–1041.
- Lupien, S. J., Parent, S., Evans, A. C., Tremblay, R. E., Zelazo, P. D., Corbo, V., et al. (2011). Larger amygdala but no change in hippocampal volume in 10-year-old children exposed to maternal depressive symptomatology since birth. *Proceedings of the National Academy of Sciences*, 108, 14324–14329.
- MacMaster, F. P., & Kusumakar, V. (2004). Hippocampal volume in early onset depression. *BMC Medicine*, 2, 2.
- MacMaster, F. P., Mirza, Y., Szeszko, P. R., Kmiecik, L. E., Easter, P. C., Taormina, S. P., et al. (2008). Amygdala and hippocampal volumes in familial early onset major depressive disorder. *Biological Psychiatry*, 63, 385–390.
- Matthews, S., Strigo, I., & Simmons, A. (2008). Decreased functional coupling of the amygdala and supragenual cingulate is related to increased depression in unmedicated individuals with current major depressive disorder. *Journal of Affective Disorders*, 111, 13–20.
- Mayberg, H. S., Brannan, S. K., Tekell, J. L., Silva, J. A., Mahurin, R. K., McGinnis, S., et al. (2000). Regional metabolic effects of fluoxetine in major depression: Serial changes and relationship to clinical response. *Biological Psychiatry*, 48, 830–843.
- McRae, K., Gross, J. J., Weber, J., Robertson, E. R., Sokol-Hessner, P., Ray, R. D., et al. (2012). The development of emotion regulation: An fMRI study of cognitive reappraisal in children, adolescents and young adults. *Social Cognitive and Affective Neuroscience*, 7, 11–22.
- Monk, C. S., Klein, R. G., Telzer, E. H., Schroth, E. A., Mannuzza, S., Moulton, J. L., et al. (2008). Amygdala and nucleus accumbens activation to

- emotional facial expressions in children and adolescents at risk for major depression. *American Journal of Psychiatry*, *165*, 90–98.
- Monk, C. S., McClure, E. B., Nelson, E. E., Zarahn, E., Bilder, R. M., Leibenluft, E., et al. (2003). Adolescent immaturity in attention-related brain engagement to emotional facial expressions. *NeuroImage*, *20*, 420–428.
- Moore, W. E., Pfeifer, J. H., Masten, C. L., Mazziotta, J. C., Iacoboni, M., & Dapretto, M. (2012). Facing puberty: Associations between pubertal development and neural responses to affective facial displays. *Social Cognitive and Affective Neuroscience*, *7*, 35–43.
- Morris, M. C., Rao, U., & Garber, J. (2012). Cortisol responses to psychosocial stress predict depression trajectories: Social–evaluative threat and prior depressive episodes as moderators. *Journal of Affective Disorders*, *143*, 223–230.
- Ochsner, K., & Gross, J. (2005). The cognitive control of emotion. *Trends in Cognitive Sciences*, *9*, 242–249.
- Ochsner, K., Ray, R., Cooper, J., Robertson, E., Chopra, S., Gabrieli, J., et al. (2004). For better or for worse: Neural systems supporting the cognitive down-and-up-regulation of negative emotion. *NeuroImage*, *23*, 483–499.
- Ochsner, K. N., Bunge, S. A., Gross, J. J., & Gabrieli, J. D. E. (2002). Rethinking feelings: An fMRI study of the cognitive regulation of emotion. *Journal of Cognitive Neuroscience*, *14*, 1215–1229.
- Ochsner, K. N., & Feldman Barrett, L. (2001). A multiprocess perspective on the neuroscience of emotion. In T. J. Mayne & G. A. Bonanno (Eds.), *Emotion: Current issues and future directions* (pp. 38–81). New York: Guilford Press.
- Ochsner, K. N., & Gross, J. J. (2007). The neural architecture of emotion regulation. In J. J. Gross (Ed.), *Handbook of emotion regulation* (Vol. 1, pp. 87–109). New York: Guilford Press.
- Ochsner, K. N., & Gross, J. J. (2008). Cognitive emotion regulation insights from social cognitive and affective neuroscience. *Current Directions in Psychological Science*, *17*, 153–158.
- Ochsner, K. N., & Phelps, E. (2007). Emerging perspectives on emotion–cognition interactions. *Trends in Cognitive Sciences*, *11*, 317–318.
- Ochsner, K. N., Ray, R. R., Hughes, B., McRae, K., Cooper, J. C., Weber, J., et al. (2009). Bottom-up and top-down processes in emotion generation: Common and distinct neural mechanisms. *Psychological Science*, *20*, 1322–1331.
- Ojemann, J. G. J., Akbudak, E. E., Snyder, A. Z. A., McKinstry, R. C. R., Raichle, M. E. M., & Conturo, T. E. T. (1997). Anatomic localization and quantitative analysis of gradient refocused echo-planar fMRI susceptibility artifacts. *NeuroImage*, *6*, 156–167.
- Ollinger, J. M., Shulman, G. L., & Corbetta, M. (2001). Separating processes within a trial in event-related functional MRI. *NeuroImage*, *13*, 210–217.
- Pagliaccio, D., Luby, J., Gaffrey, M., Belden, A., Botteron, K., Gotlib, I. H., et al. (2011). Anomalous functional brain activation following negative mood induction in children with pre-school onset major depression. *Developmental Cognitive Neuroscience*, *2*, 256–267.
- Pagliaccio, D., Luby, J. L., Gaffrey, M. S., Belden, A. C., Botteron, K. N., Harms, M. P., et al. (2013). Functional brain activation to emotional and nonemotional faces in healthy children: Evidence for developmentally undifferentiated amygdala function during the school-age period. *Cognitive, Affective, & Behavioral Neuroscience*, *13*, 771–789.
- Park, H. D., & Tallon-Baudry, C. (2014). The neural subjective frame: From bodily signals to perceptual consciousness. *Philosophical Transactions of the Royal Society B: Biological Sciences*, *369*, 20130208.
- Peng, J., Liu, J., Nie, B., Li, Y., Shan, B., Wang, G., et al. (2011). Cerebral and cerebellar gray matter reduction in first-episode patients with major depressive disorder: A voxel-based morphometry study. *European Journal of Radiology*, *80*, 395–399.
- Power, J. D., Barnes, K. A., Snyder, A. Z., Schlaggar, B. L., & Petersen, S. E. (2013). Steps toward optimizing motion artifact removal in functional connectivity MRI: A reply to Carp. *NeuroImage*, *76*, 439–441.
- Power, J. D., Mitra, A., Laumann, T. O., Snyder, A. Z., Schlaggar, B. L., & Petersen, S. E. (2013). Methods to detect, characterize, and remove motion artifact in resting state fMRI. *NeuroImage*. Advance online publication.
- Rao, U., Chen, L.-A., Bidesi, A. S., Shad, M. U., Thomas, M. A., & Hammen, C. L. (2010). Hippocampal changes associated with early-life adversity and vulnerability to depression. *Biological Psychiatry*, *67*, 357–364.
- Roberson-Nay, R., McClure, E. B., Monk, C. S., Nelson, E. E., Guyer, A. E., Fromm, S. J., et al. (2006). Increased amygdala activity during successful memory encoding in adolescent major depressive disorder: An fMRI study. *Biological Psychiatry*, *60*, 966–973.
- Rosso, I. M., Cintron, C. M., Steingard, R. J., Renshaw, P. F., Young, A. D., & Yurgelun-Todd, D. A. (2005). Amygdala and hippocampus volumes in pediatric major depression. *Biological Psychiatry*, *57*, 21–26.
- Roy, A. K., Shehzad, Z., Margulies, D. S., Kelly, A. M. C., Uddin, L. Q., Gotimer, K., et al. (2009). Functional connectivity of the human amygdala using resting state fMRI. *NeuroImage*, *45*, 614–626.
- Saleh, K., Carballo, A., Liseicka, D., Fagan, A. J., Connolly, G., Boyle, G., et al. (2012). Impact of family history and depression on amygdala volume. *Psychiatry Research*, *203*, 24–30.
- Sauder, C. L., Hajcak, G., Angstadt, M., & Phan, K. L. (2013). Test–retest reliability of amygdala response to emotional faces. *Psychophysiology*, *50*, 1147–1156.
- Sheline, Y. I., Gado, M. H., & Price, J. L. (1998). Amygdala core nuclei volumes are decreased in recurrent major depression. *NeuroReport*, *9*, 2023–2028.
- Sheline, Y. I., Price, J. L., Yan, Z., & Mintun, M. A. (2010). Resting-state functional MRI in depression unmasks increased connectivity between networks via the dorsal nexus. *Proceedings of the National Academy of Sciences*, *107*, 11020–11025.
- Sheline, Y. I., Sanghavi, M., Mintun, M. A., & Gado, M. H. (1999). Depression duration but not age predicts hippocampal volume loss in medically healthy women with recurrent major depression. *Journal of Neuroscience*, *19*, 5034–5043.
- Siegle, G., Carter, C., & Thase, M. (2006). Use of fMRI to predict recovery from unipolar depression with cognitive behavior therapy. *American Journal of Psychiatry*, *163*, 735.
- Siegle, G. J., Konecky, R. O., Thase, M. E., & Carter, C. S. (2003). Relationships between amygdala volume and activity during emotional information processing tasks in depressed and never-depressed individuals: An fMRI investigation. *Annals of the New York Academy of Sciences*, *985*, 481–484.
- Silk, J. S., Steinberg, L., & Morris, A. S. (2003). Adolescents' emotion regulation in daily life: Links to depressive symptoms and problem behavior. *Child Development*, *74*, 1869–1880.
- Silvers, J. A., McRae, K., Gabrieli, J. D. E., Gross, J. J., Remy, K. A., & Ochsner, K. N. (2012). Age-related differences in emotional reactivity, regulation, and rejection sensitivity in adolescence. *Emotion*, *12*, 1235–1247.
- Stein, J. L., Wiedholz, L. M., Bassett, D. S., Weinberger, D. R., Zink, C. F., Mattay, V. S., et al. (2007). A validated network of effective amygdala connectivity. *NeuroImage*, *36*, 736–745.
- Stein, M. B., Koverola, C., Hanna, C., Torchia, M. G., & McClarty, B. (1997). Hippocampal volume in women victimized by childhood sexual abuse. *Psychological Medicine*, *27*, 951–959.
- Stuhrmann, A., Suslow, T., & Dannlowski, U. (2011). Facial emotion processing in major depression: A systematic review of neuroimaging findings. *Biology of Mood and Anxiety Disorders*, *1*, 10.
- Surguladze, S., Brammer, M. J., Keedwell, P., Giampietro, V., Young, A. W., Travis, M. J., et al. (2005). A differential pattern of neural response toward sad versus happy facial expressions in major depressive disorder. *Biological Psychiatry*, *57*, 201–209.
- Surguladze, S. A., Brammer, M. J., Young, A. W., Andrew, C., Travis, M. J., Williams, S. C. R., et al. (2003). A preferential increase in the extrastriate response to signals of danger. *NeuroImage*, *19*, 1317–1328.
- Suzuki, H., Botteron, K. N., Luby, J. L., Belden, A. C., Gaffrey, M. S., Babb, C. M., et al. (2012). Structural–functional correlations between hippocampal volume and cortico-limbic emotional responses in depressed children. *Cognitive, Affective, & Behavioral Neuroscience*, *13*, 135–151.
- Talairach, J., & Tournoux, P. (1988). *Co-planar stereotaxic atlas of the human brain. 3-Dimensional proportional system: An approach to cerebral imaging*. New York: Thieme.
- Taylor, L., & Ingram, R. E. (1999). Cognitive reactivity and depressotypic information processing in children of depressed mothers. *Journal of Abnormal Psychology*, *108*, 202–210.
- Thomas, K., Drevets, W., Whalen, P., Eccard, C., Dahl, R., Ryan, N., et al. (2001). Amygdala response to facial expressions in children and adults. *Biological Psychiatry*, *49*, 309–316.
- Tottenham, N., Hare, T. A., Quinn, B. T., McCarry, T. W., Nurse, M., Gilhooly, T., et al. (2010). Prolonged institutional rearing is associated with atypically large amygdala volume and difficulties in emotion regulation. *Developmental Science*, *13*, 46–61.
- Tottenham, N., Tanaka, J. W., Leon, A. C., McCarry, T., Nurse, M., Hare, T. A., et al. (2009). The NimStim set of facial expressions: Judgments from untrained research participants. *Psychiatry Research: Neuroimaging*, *168*, 242–249.
- van der Plas, E. A. A., Boes, A. D., Wemmie, J. A., Tranel, D., & Nopoulos, P. (2010). Amygdala volume correlates positively with fearfulness in normal healthy girls. *Social Cognitive and Affective Neuroscience*, *5*, 424–431.

- Van Eijndhoven, P., van Wingen, G., van Oijen, K., Rijpkema, M., Goraj, B., Verkes, J. R., et al. (2009). Amygdala volume marks the acute state in the early course of depression. *Biological Psychiatry*, *65*, 812–818.
- Vassilopoulou, K., Papathanasiou, M., Michopoulos, I., Boufidou, F., Oulis, P., Kelekis, N., et al. (2012). A magnetic resonance imaging study of hippocampal, amygdala and subgenual prefrontal cortex volumes in major depression subtypes: Melancholic versus psychotic depression. *Journal of Affective Disorders* *146*, 197–204.
- Videbech, P., & Ravnkilde, B. (2004). Hippocampal volume and depression: A meta-analysis of MRI studies. *American Journal of Psychiatry*, *161*, 1957–1966.
- Vink, M., Derks, J. M., Hoogendam, J. M., Hillegers, M., & Kahn, R. S. (2014). Functional differences in emotion processing during adolescence and early adulthood. *NeuroImage*, *91*, 70–76.
- Westermann, R., Spies, K., Stahl, G., & Hesse, F. W. (1996). Relative effectiveness and validity of mood induction procedures: A meta-analysis. *European Journal of Social Psychology*, *26*, 557–580.
- Whittle, S., Simmons, J. G., Dennison, M., Vijayakumar, N., Schwartz, O., Yap, M. B. H., et al. (2013). Positive parenting predicts the development of adolescent brain structure: A longitudinal study. *Developmental Cognitive Neuroscience*, *8*, 7–17.
- Williams, L. M., Phillips, M. L., Brammer, M. J., Skerrett, D., Lagopoulos, J., Rennie, C., et al. (2001). Arousal dissociates amygdala and hippocampal fear responses: Evidence from simultaneous fMRI and skin conductance recording. *NeuroImage*, *14*, 1070–1079.
- Woon, F. L., & Hedges, D. W. (2008). Hippocampal and amygdala volumes in children and adults with childhood maltreatment-related posttraumatic stress disorder: A meta-analysis. *Hippocampus*, *18*, 729–736.
- Yang, T. T., Simmons, A. N., Matthews, S. C., Tapert, S. F., Frank, G. K., Max, J. E., et al. (2010). Adolescents with major depression demonstrate increased amygdala activation. *Journal of the American Academy of Child & Adolescent Psychiatry*, *49*, 42–51.
- Zeman, J., Shipman, K., & Penza-Clyve, S. (2001). Development and initial validation of the Children's Sadness Management Scale. *Journal of Nonverbal Behavior*, *25*, 187–205.